

Case No.: SJKIM-002US
Patent Application
Express Mail

**COMPOSITION CONTAINING ASIASARI RADIX EXTRACTS
FOR PROTECTING BRAIN CELLS AND IMPROVING MEMORY**

BACKGROUND OF THE INVENTION

[0001] This invention is related to a composition for protecting brain cells and improving memory comprising Asiasari Radix extracts.

[0002] One of the major factors involved in brain cell damages is glutamate. Glutamate exerts its actions by binding to four types of receptors including NMDA receptor, AMPA receptor, Kainate receptor and 1S,3R-ACPD receptor [Craig CR, Stitzel RE, Modern Pharmacology with clinical applications, 293-302, 1997]. When brain ischemia occurs, oxygen supply to brain cells diminishes and then anaerobic glycolysis increases. Therefore, ATP content in the cell decreases and the concentration of extracellular potassium ions increases. Eventually, depolarization of neuronal cell membrane and subsequent release of excitatory amino acids occurs which results in the neuronal damages by the activation of receptors for NMDA, AMPA and Kainate. Excitotoxicity via excitatory neurotransmitters induces cellular stress and thereby plays an important role in the pathogenesis of neurodegenerative disorders such as Alzheimer's disease, Parkinson disease, strokes and amyotrophic lateral sclerosis [Halloween, B., J. Neurochem. 59,

1609-1623, 1992; Coyle, J. T. and Puttfarcken, P., Science 262, 689-695. 1993; Olanow, C. W., Trends. Neurosci. 16, 439-444, 1993].

[0003] Neurodegenerative disorders in the central nervous system are often accompanied by a decrease in cognition and memory. Especially, dementia is a serious problem of modern societies with high population of elderly. Dementia is typically caused by a variety of environmental factors such as genetics, aging, brain damage, smoking and alcohol and other complex factors. The hippocampus of patients suffering from dementia is heavily damaged and this is closely related to the reduction of acetylcholine levels in the brain.

[0004] Acetylcholinesterase inhibitors are clinically used for the Alzheimer's disease to increase acetylcholine levels. Recently, a great number of researches have been performed to search for ways of neuroprotection such as by using NMDA antagonist, AMPA antagonist, GABA agonist, intracellular calcium decreasing agent, nitric oxide inhibitor, free radical scavenger, Na channel blocker, inhibitor of glutamate release, acidosis, hypothermia and potassium channel activator [Gagliardi RJ, Neuroprotection, excitatotoxicity and NMDA antagonists, Arq. Neuro-Psiquiatr. 58, 2000].

[0005] Dozocylpin (MK 801), selfotel, cerestat, and dextrometorfan have been developed as NMDA antagonists. However, they are known to induce altered sensory perception, dysphoria, nystagmus, and hypotension at low doses, while inducing psychological adverse events such as excitement, paranoia, and hallucination at higher doses. Furthermore, NBQX was developed as

an AMPA antagonist. However, it causes severe kidney toxicity and is thus not optimal in clinical applications. Therefore, neuroprotective agents without toxicity from natural product-derived materials need to be developed.

[0006] Recently, it has been found that AMPA receptor plays an important role in the development of Alzheimer's disease. The fact that neuronal cell damages by the AMPA receptor activation occurs selectively at the basal forebrain cholinergic neurons (BFCNs) suggests that development of anti-Alzheimer's disease can be tried by using AMPA receptor antagonists [Weiss, J. H., et al., Basal forebrain cholinergic neurons are selectively vulnerable to AMPA/kainate receptor-mediated neurotoxicity. Neuroscience 60, 659-664, 1994].

[0007] Glial cells are essential for the survival of neuronal cells. In developing the central nervous system, glial cells control the precise movement and growth of neuronal cells whereas they play a role in homeostasis and synaptic plasticity of neuronal cells after development. In addition, glial cells contain receptors and neurotransmitters capable of initiating neuronal signal transduction essential for survival and apoptosis of neuronal cells. Thus, protecting glial cells from external damages are eventually related to the plasticity, homeostasis and survival of neuronal cells.

[0008] Insulin receptor in the peripheral tissues participates mainly in the glucose metabolism whereas its role in the CNS appeared not to be related to glucose metabolism but other neuronal

activity such as memory. In fact, insulin receptor is widely distributed in different areas in the brain and present in a large amount in the hippocampus. Hence, the hippocampus is an important target of insulin action in the brain. Recently, much evidence has been presented as to the role of brain insulin or insulin receptors in the memory formation. It has been found that both experimental damage to the neuronal insulin receptor and Alzheimer brain induced similar metabolic abnormalities [Hoyer, S., Muller, D, Plaschke, K. Desensitization of brain insulin receptor. Effect on glucose/energy and related metabolism. J. Neural Transm [Suppl] 44, 259-268, 1994].

[0009] Interesting hypothesis has been proposed that sporadic Alzheimer disease might be the brain type of non-insulin dependent diabetes mellitus (Hoyer, S. Is sporadic Alzheimer disease the brain type of non-insulin dependent diabetes mellitus? A challenging hypothesis. J. Neural Transm. 105, 415-422, 1998). It has been suggested that intracerebroventricular insulin enhances memory in a passive-avoidance task [Park, C. P., Seeley, R. J., Craft, S. and Woods S. C. (2000) Intracerebroventricular insulin enhances memory in a passive avoidance task. Physiol. Behav. 68, 509-514]. Insulin receptor density and tyrosine kinase activity in the sporadic Alzheimer's disease (SDAT) was known to be significantly decreased [Frolich, L., Blum-degen, D., Bernstein, H. G., Engelsberger, S., Humrich, J., Laufer, S., Muschner, D., Thalheimer, A., Turk, A., Hoyer, S., Zochling, R., Boissl, K. W., Jellinger, K., and Piederer, P. Brain insulin and insulin receptors

in aging and sporadic Alzheimer's disease. J. Neural Transm. 105, 423-438, 1998]. Interestingly, tyrosine phosphorylation of the hippocampal insulin receptor has been shown to play an essential role in spatial memory formation [Zhao, W., Chen, H., Xu, H., Moore, E., Meiri, N., Quon, M. J., Alkon, D. L. (1999) Brain insulin receptors and spatial memory. J. Biol. Chem. 274, 34893-34902, 1999]. Taken together, insulin receptor activators could be used for memory enhancement in addition to cholinesterase inhibitors.

BRIEF SUMMARY OF THE INVENTION

[0010] The present invention is related to a composition containing Asiasari Radix extracts for protecting brain cells and improving memory. It can be used as drugs and/or health foods for the prevention and treatment of neurodegenerative diseases via brain cell protection of the moderns who are being subjected to brain damages due to various kinds of stress, drinking, smoking, etc., as well as for the improvement of memory.

BRIEF DESCRIPTION OF THE DRAWINGS

[0011] These as well as other features of the present invention will become more apparent upon reference to the drawings wherein:

[0012] Fig. 1 shows fraction 1 of Asiasari Radix (AR) extract blocking the depolarization induced by AMPA in the rat cortical slices. Data are expressed as mean SD (n=5). *: p 0.05 with respect to the control;

[0013] Fig. 2 shows fraction 4 of AR extract blocking the depolarization induced by AMPA in the rat cortical slices. Data are expressed as mean SD (n=5). **: p 0.01 with respect to the control;

[0014] Fig. 3 shows fraction 1 of AR extract having inhibitory actions against neuronal cell damages induced by AMPA in the differentiated PC12 cells. Data are expressed as mean SD (n=5). ***: p 0.001 with respect to the control;

[0015] Fig. 4 shows the effects of fractions 2, 3, 4 and 5 of AR extract on neuronal cell damages induced by AMPA in the differentiated PC12 cells. Data are expressed as mean SD (n=5). ***: p 0.001 with respect to the control;

[0016] Fig. 5 shows fraction 1 of AR extract having inhibitory actions against cell damages induced by AMPA in the C6 glial cells. Data are expressed as mean SD (n=5). Bars not sharing a common letter are different at p 0.05 by Duncan's multiple range test;

[0017] Fig. 6 shows fractions 1, 2, 3, 4 and 5 of AR extract having inhibitory actions against glial cell damages induced by ZnCl₂ in the C6 glial cells. Data are expressed as mean SD (n=5). *: p 0.05, ***: p 0.001 with respect to the control;

[0018] Fig. 7 shows fraction 1 of AR extract having increased memory in the NaNO₂ assay. Data are expressed as mean SD (n=8). **: p 0.01 with respect to the control;

[0019] Fig. 8 shows the effects of fractions 2, 3, 4 and 5 of AR extract on the memory in the NaNO₂ assay. Data are expressed as mean SD (n=8). *: p 0.05, **: p 0.01 with respect to the control;

[0020] Fig. 9 shows administration of fraction 2 of AR extract via P.O. causing a significant increase of memory in the NaNO_2 assay. Data are expressed as mean SD (n=8). *: p 0.05 with respect to the control;

[0021] Fig. 10 shows the fractions 1, 2 and 4 of AR extract stimulating memory enhancement in the 8 radial arm maze assay. Data are expressed as mean SD (n=10). *: p 0.05, **: p 0.01 with respect to the control;

[0022] Fig. 11. shows fractions 1, 2 and 4 of AR extract stimulating memory enhancement in the passive avoidance test. Data are expressed as mean SD (n=10). *: p 0.05 with respect to the control;

[0023] Fig. 12 shows fractions 1, 2 and 4 of AR extract stimulating tyrosine phosphorylation of rat hippocampal proteins.;

[0024] Fig. 13 shows tyrosine phosphorylation of insulin receptors by fractions 1, 2 and 4 of AR extract in rat hippocampus;

[0025] Fig. 14 shows fractions 1, 2 and 4 of AR extract activating ERK I and ERK II in rat hippocampus; and

[0026] Fig. 15 shows fractions 1, 2 and 4 of AR extract inhibiting cholinesterase activity in rat hippocampus.

DETAILED DESCRIPTION OF THE INVENTION

[0027] The inventor of the present invention has studied for many years to search for neuroprotective agents with memory enhancing activity for the moderns who are being subjected to brain damages due to environmental factors such as various stress,

alcohol, smoking etc. Based on the search of many natural products, it was found that Asiasari Radix extracts have significant neuroprotective effects as well as memory enhancing activity.

[0028] In this respect, the present invention is related to a composition containing Asiasari Radix extracts for protecting brain cells and improving memory. A feature of the composition of the present invention is that it contains 0.5 to 50% by weight of Asiasari radix extracts per total weight of the composition.

[0029] Asiasari Radix is the dried wholeplant, rhizome or root of *Asarum sieboldi* F. MAEKAWA, *Asarum sieboldi* var. *seoulense* Nakai, *Asarum sieboldii* Miq., *Asarum heterotropoides* Schmidt, *Asarum heterotropoides* var. *mandshuricum* Kitagawa, or *Asiasarum heterotropoides* F. MAEKAWA var. *seoulensis* F. MAEKAWA. It is a perennial plant. Its rhizome projecting aside is relatively short and it contains a lot of knobs and fine roots with a diameter of approximately 1 mm. It is shaped like an uneven and crooked string. It has a knotty rhizome with a diameter of 3-5 mm and is yellowish brown. The rhizome contains a lot of roots with 5-20 cm in length. It has a light brown color and is long and slender. It has very shallow vertical pleats.

[0030] It contains essential oils such as methyleugenol, asarylketone, cineol, safrole, limonen, eucarvone and acidamide such as N-isobutyl 2, 4, 8, 10-dodecatetraenamide, pellitorin and lignans such as sesamin and asarinin (Zhou RH, Resource Science of Chinese Medicinal Materials. pp.202-211. Beijing: China medical &

Pharmaceutical Sciences press, 1993). Several alkaloids including hygenamine are also present in Asiasari Radix. Extracts of Asiasari Radix have been reported to exert body temperature decreasing, spasmolytic, antihistamine and cardiogenic actions. It is used as local anesthetics, fever remedy, cough remedy, expectorant and diuresis. (Zhu, Y. Chinese Materia Medica: Chemistry, Pharmacology and Application, pp. 66-69. Beijing: People's Health Publisher, 1998). However, it is not known as to whether Asiasari Radix extract has any effects on neuroprotection and memory.

[0031] Asiasari Radix extracts of the present invention are obtained by extracting Asiasari Radix with a lower alcohol having 1 carbon atom to 4 carbon atoms such as methanol or ethanol, and organic solvent such as acetone, chloroform, methylenechloride, ether or ethylacetate at 5°C to 80°C, desirably 30°C to 55°C for 15 min to 48hrs, desirably 30 min to 12 hrs. The obtained extracts can be made as powders by evaporation under reduced pressure.

[0032] In addition, the Asiasari Radix extracts of the present invention could be further fractionated by the following method (Harborne J. B. Phytochemical methods: A guide to modern techniques of plant analysis. 3rd Edt. Pp 6-7, 1998).

[0033] The Asiasari Radix extracts of the present invention are prepared by the following sequential extraction and fractionation procedure: the extracts of Asiasari Radix obtained by the above mentioned method are dissolved in a methanol:water mixed solvent, adjusted to pH 2-4 and followed by extraction with equal volume of

chloroform; the chloroform-insoluble fraction is adjusted to pH 9-12 with NH₄OH and subjected to extraction with equal volume of a chloroform:methanol mixed solvent; among these, a fraction insoluble in the chloroform:methanol mixed solvent is subsequently subjected to extraction with methanol to obtain the Asiasari Radix extracts.

[0034] Desirable ratios of chloroform:methanol mixture solvent are in the ranges of 1: 0.1~1. When the chloroform-insoluble fractions described above are further fractionated with chloroform:methanol, the fraction soluble in the chloroform:methanol mixed solvent contains most alkaloids while the fraction insoluble in the chloroform:methanol mixture solvent contains quaternary alkaloids and N-oxides.

[0035] The composition comprising Asiasari Radix extracts could contain suitable carriers, excipients and diluents such as lactose, dextrose, sucrose, sorbitol, mannitol, xylitol, erythritol, maltitol, starch, acacia gum, alginate, gelatin, calcium, phosphate, calcium silicate, cellulose, methyl cellulose, microcrystalline cellulose, polyvinyl pyrrolidone, water, methylhydroxy-benzoate, propylhydroxybenzoate, talc, magnesium stearate or mineral oil. The composition according to the present invention can be used as oral formulations such as powders, tablets, capsules, suspensions, emulsions, syrups, aerosol and external application and suppositories. In addition, it could be used as sterilized injections. The Asiasari Radix extracts in the present invention can be administered at the dosage of 0.1 mg/Kg to

500 mg/Kg per day. The dosage can be administered once or it can be administered as divided. However, the actual dose administered should be understood in terms of various related factors such as kind of diseases to treat, administration route, age of patient, gender and weight, and the condition of disease, etc. Therefore, above-mentioned dosage does not limit the range of the present invention by any ways. The composition comprising Asiasari Radix extracts in the present invention can be widely used as drugs, foods and drinks for neuroprotection and memory improvement. Asiasari Radix extracts can be added to foods such as various foods, drinks, gum, tea, vitamin complexes and health foods etc.

[0036] In addition, the present invention is related to a composition for improving memory containing a chloroform fraction of Asiasari Radix extracts obtained by the following sequential fractionation procedure: Asiasari Radix is subjected to extraction with a lower alcohol having between 1 carbon atom and 4 carbon atoms such as methanol or ethanol, or organic solvent such as acetone, chloroform, methylene chloride, ether or ethylacetate; the resulting Asiasari Radix extracts are solubilized in a methanol:water mixed solvent, adjusted to pH 2-4 with acid and subjected to extraction with equal volume of chloroform to obtain the chloroform fraction of Asiasari Radix extracts. Desirable ratios of the methanol:water mixed solvent are in the ranges of 1:0.2~1.5 and this fraction contains terpenoids and phenolic compounds.

[0037] The composition for improving memory of the present

invention has a feature that the content of the fraction described above ranges 0.5 to 50 % by weight per total weight of the composition.

[0038] The composition for improving memory of the present invention could contain suitable carriers, excipients and diluents as same as the composition containing Asiasari Radix extracts for protecting brain cells and improving memory as described above. In addition, it could be made as various formulations according to conventional methods and be widely used as drugs, foods, and drinks for improving memory.

[0039] The present invention is explained in detail by the following examples, but it is not limited to these examples.

[0040] **Example: Preparation of Asiasari Radix extracts**

Asiasari Radix (250 g) was cut into small pieces and extracted with 70% methanol (750 ml) by the use of Soxhlet apparatus for 3 hours three times. Following filtration, the resulting methanol extract was concentrated by rotary evaporator and dried by freeze-dryer (Fr. 1). For fractionation with other organic solvents, 10g of Fr. 1 was resuspended with 200 ml MeOH-H₂O (4:1), which was then adjusted to pH 3 with 2M H₂SO₄. Then, it was extracted with equal volume of CHCl₃ three times and concentrated using a rotary evaporator, followed by freeze-dried (Fr. 2).

[0041] The water-soluble and CHCl₃ insoluble fraction was adjusted to pH 10 with NH₄OH and subjected to extraction with equal volume of CHCl₃-MeOH (3:1) two times. The CHCl₃-MeOH (3:1) soluble fraction was concentrated and freeze dried (Fr. 3). The water-

soluble and CHCl_3 -MeOH (3:1) insoluble fraction was mixed with equal volume of MeOH and subjected to extraction three times. The MeOH soluble fraction was concentrated and freeze-dried (Fr. 4). The MeOH insoluble fraction was concentrated and freeze-dried (Fr. 5).

[0042] Experiment 1. Grease-gap recording assay

[0043] 1) Experimental methods

[0044] Cortical wedges from rats were prepared, placed in two-compartment brain bath and experiments were performed as described by Harrison and Simmonds (British J. Pharmacol. 84, 381-391, 1985). 'Wedges' of rat cerebral cortex were prepared from the brains of male Wistar rats (200 g). The animal is decapitated and the brain rapidly removed into chilled oxygenated (95% O_2 / 5% CO_2) artificial cerebrospinal fluid (ACSF) containing (mM): NaCl 122, NaHCO_3 25, KCl 3.1, KH_2PO_4 0.4, CaCl_2 1.3, MgSO_4 1.4, D-glucose 10, (pH 7.4). 500-600 μ m coronal slices were taken and they were then placed in room temperature oxygenated ACSF and hemisected with a razor blade. A wedge of tissue was formed such that the dorsal cortical surface containing cerebral cortex and corpus callosum was approximately 1.5 mm wide and ventral surface was approximately 1mm wide. The wedges were further incubated in ACSF for 2 hrs at room temperature.

[0045] Cortical wedges were placed in a two-compartment bath and a greased (high vacuum silicon grease) barrier placed such that a high-resistance seal was formed between the two compartments. Oxygenated ACSF was perfused through the two compartments

separately at 2 ml/min for at least 1 hr. Asiasari Radix (AR) extracts (fractions 1, 2, 3 or 4) were perfused 10 min before the application of AMPA (40 M) for 2 min: AR extracts and AMPA were applied to the cortical compartment. The DC potential between the two compartments was monitored via Ag/AgCl electrodes. The signal was amplified and analyzed with the aid of McLab software.

[0046] 2) Experimental results

[0047] Depolarization of neuronal cells by AMPA is regarded as an index of stimulation by damages of neuronal cells. AMPA treatment (40 uM) for 2 min caused an increase in depolarization up to 0.79 mV. When AR extract (fraction 1) (10 ug/ml) was preincubated for 10 min and followed by AMPA treatment for 2 min, AMPA-induced increase in the depolarization was significantly decreased (Fig. 1A). The average inhibitory effect by the fraction 1 of AR extract was about 54% as compared to the control (Fig. 1B). These results suggest that AR extract (fraction 1) has significant neuroprotective actions against AMPA-induced neuronal damages.

[0048] Among the fractions from methanol extract of Asiasari Radix, fraction 4 has dramatic inhibitory action against AMPA-induced depolarization: it was about 92% protection against AMPA-induced neuronal damages (Fig. 2). On the other hand, fractions 2 and 3 were ineffective in the inhibition against AMPA-induced depolarization (data not shown).

[0049] Since the fraction 4 contains components which are insoluble in chloroform as well as in chloroform:methanol, it is devoid of terpenoids, phenolic compounds and sesamin which are

soluble in organic solvents such as chloroform. Thus, it is suggested that the neuroprotective effects of fraction 4 be brought about by components other than sesamin, terpenoid and phenolic compounds etc.

[0050] Experiment 2. Cell viability assay (MTT assay)

[0051] 1) Experimental methods

[0052] MTT assay is a method to measure mitochondrial redox with colorimeter and it is primarily used to know mitochondrial redox potential or cell viability (Mosmann T., J Immunol. Methods, 1983). PC 12 cells and C6 glial cells were grown in DMEM media with 10% fetal calf serum in 5% humidified CO₂ atmosphere at 37°C. PC12 cells were induced to differentiation into neuronal cells by the addition of nerve growth factor (100 ng/ml). Fresh NGF was added every 48 h and experiments were conducted after NGF treatment for 7 days.

[0053] The cells were plated in 96 well plates at a density of 1×10^5 cells/well. Cells were preincubated with AMPA (40 M) and followed by the addition of AR fractions (10 g/ml) for 24 hrs. MTT reagent (Sigma, USA) (5 mg/ml) was made in PBS (phosphate buffered saline) and filtered. The cells were then treated with MTT (final concentration, 0.5 mg/ml) and allowed to incubate for 3 hrs at 37°C. Cells containing active mitochondria form dark blue formazan by disintegrating tetrazolium ring. The culture media were removed and the cells were subjected to lysis in the presence of 100 μ l of DMSO and 10 μ l of Sorensen glycine buffer (0.1M glycine, 0.1M NaCl, pH 10.5). The absorbance was measured with spectrophotometer at 570

nm. The ratio of absorbance of experimental groups to control groups is expressed as % cell viability.

[0054] 2) Experimental results

[0055] In the differentiated neuronal PC12 cells, AMPA (40 uM) caused approximately 50% reduction in the cell viability as compared to control whereas AR extract (fraction 1; 10 ug/ml) pretreatment caused more than 90% recovery from the reduction of cell viability caused by AMPA (Fig. 3). When fractions 2, 3, 4 or 5 were tested for their protective actions on cell viability, fraction 4 protected the AMPA-induced neuronal damages in the differentiated PC12 cells almost to the control level, suggesting it has active principles responsible for neuroprotection (Fig. 4). In C6 glial cells, AMPA caused approximately 35% reduction in the cell viability as compared to control whereas AR extract (fraction 1) pretreatment caused approximately 10% recovery from the reduction of cell viability caused by AMPA (Fig. 5). Since the original AR extract has relatively weaker protective effects against glial cell damages induced by AMPA, we tried to induce oxidative damages in the C6 glial cells with ZnCl₂ (100 uM) (Fig. 6). When fractions 1, 2, 3, 4 or 5 were pretreated, fraction 1 blocked the glial cell damages by 100% induced by Zn and fraction 4 inhibited the damages by 58% as compared to control.

[0056] Experiment 3. Memory assay

[0057] 1. NaNO₂ assay

[0058] It has been suggested that there is close relationship between deficit of oxidative metabolism in the brain by NaNO₂ and

cholinergic neurotransmission related to learning and memory. Therefore, the prolongation of survival time is regarded as an index of memory enhancement [Schindler et al., Drug Develop. Res. 4: 567-576, 1984].

[0059] 1) Experimental methods

[0060] Asiasari radix extract (fraction 1; 100 mg/Kg, I. P.) were administered to male mouse. After 60 min, NaNO_2 (250 mg/Kg, S. C.) was injected and cessation of respiration was recorded.

The memory enhancing effects were analyzed by measuring and comparing the prolongation of survival times of control and treated groups.

[0061] The subfractions (fraction 2, 3, 4, and 5) of fraction 1 were also administered to male mouse (100 mg/kg, i.p.) and tested for the NaNO_2 memory assay as described above.

[0062] 2) Experimental results

[0063] The original methanol extract of Asiasari Radix (fraction 1) administration (100 mg/Kg, I. P.) caused a 25% increase in the survival time as compared to control, indicating that it improves memory (Fig. 7). When fractions 2, 3, 4 and 5 were tested, fraction 4 caused about 25% increase in the survival time whereas fraction 2 caused the strongest memory enhancing effects (50% increase) among the fractions tested (Fig. 8). Subsequently, the fraction 2 was further tested as to whether oral administration could cause similar effects as i.p. administration. Fraction 2 was administered orally (10 mg/Kg) and after 60 min NaNO_2 was administered. Administration of fraction 2 (10 mg/Kg, P. O.)

caused approximately 50% increase in the memory (Fig. 9).

[0064] 2. 8 arm radial maze test

[0065] To further improve that the fractions 1, 2 and 4 have memory enhancing activity, the 8 arm radial arm maze test was carried out as described by Ikonen and Riekkinen [European J Pharmacol. 382, 151-156, 1999].

[0066] 1) Experimental methods

[0067] Central part of the 8 arm radial maze measured 20 cm in diameter. Its arms are 25 cm long, 15 cm height, 6 cm wide. Briefly, two days of pre-training (5 min per day) allowed the mice to explore the baited radial arm maze. During the experimental phase, four of the arms were baited in a semi-random manner, with a unique combination of baited arms for each mouse. After each return to the center from an arm, the doors were closed for 5 s. The training was continued until all the baits were consumed or 15 min had passed. The test involved two parameters of memory function: 1) reference memory error, entry into unbaited arms; 2) working memory error, repeated entry into arms that had been visited within a trial [Gamoh et al., Clinical and experimental Pharmacology and Physiology 28, 266-270, 2001]. The fractions 1, 2 and 4 (10 mg/Kg/day, P.O.) was administered once a day for five days and tested for the 8 arm radial maze.

[0068] 2) Experimental results

[0069] The reference memory error was reduced by 38%, 50% and 50% as compared to the control in response to fractions 1, 2 and 4, respectively while working memory error was reduced by 48%, 63% and

74% as compared to the control in response to fractions 1, 2 and 4, respectively (Fig. 10). The time taken to acquire all baits was reduced by 55%, 61% and 58% as compared to control in response to fractions 1, 2 and 4, respectively.

[0070] 3. Passive Avoidance Test

[0071] 1) Experimental methods

[0072] The test was basically performed according to the step-through method described by Jarvik and Kopp [Jarvik, M. E. and Kopp, R. An improved one-trial passive avoidance learning situation. Psychol. Rep. 21, 221-224, 1967]. The Gemini Avoidance System (SD Instruments) was used for this experiments. The apparatus consists of a two-compartment acrylic box with a lightened compartment connected to a darkened one by an automatic guillotine door. Mice were placed in the lighted box for 300 sec. Then, the guillotine door was open. Mice, as soon as they entered the dark compartment, received a punishing electrical shock (0.3 mA, 1 sec). The latency times for entering the dark compartment were measured in the training test and after 24 hr in the retention test. The maximum entry latency allowed in the retention session was 500 sec.

[0073] 3) Experimental Results

[0074] There were no significant difference among the fractions of AR extract in the training session. In the retention session, however, fraction 1, fraction 2 or fraction 4 administration caused an increase in the step-through time by 8.1-fold, 2.2-fold and 2.7-fold, respectively as compared to the control (Fig. 11).

[0075] **Experiment 4. Tyrosine phosphorylation of hippocampal proteins**

[0076] 1) **Experimental Methods**

[0077] 1. **Preparation of hippocampal lysate**

[0078] Male Sprague Dawley rats were decapitated and subjected to the isolation of hippocampus on 4°C. Hippocampal homogenates were prepared as described earlier with some modifications [Zhao, W., Chen, H., Xu, H., Moore, E., Meiri, N., Quon, M. J., Alkon, D. L., Insulin receptors and spatial memory. J. Biol. Chem. 274, 34893-34902, 1999]. The isolated hippocampus was resuspended with buffer A containing 50 mM Tris HCl, pH 7.4, 1 mM EDTA, 1 mM EGTA, 150 mM NaCl, 1% Triton X-100, 0.5 mM PMSF, 1 mM Na₃VO₄, 1 µg/ml of leupeptin and aprotinin and subjected to homogenization with a Potter-Elvehjem homogenizer. The lysates were then spun at 1,000 x g for 5 min and the supernatant were subjected to protein assay and saved at 70°C.

[0079] 2. **Immunoprecipitation**

[0080] Immunoprecipitation was performed as described earlier [Kim S. J., Kahn, C. R. Insulin stimulates phosphorylation of c-Jun, c-Fos and Fos-related proteins in cultured adipocytes. J. Biol. Chem. 269, 11887-11892, 1994]

[0081] Equal amount of proteins from hippocampal lysates were allowed to incubate with insulin receptor antibody for 1 hr at 4°C, followed by the addition of Protein A-Sepharose, and the immune complex was precipitated by centrifugation. The pellets were washed successively with 1 ml of buffer A (0.01M Tris, pH 7.4, 1M NaCl, 1%

Nonidet P-40), buffer B (0.01M Tris, pH 7.4, 0.1M NaCl, 0.01M EDTA, 1% Nonidet P-40, 0.3% SDS) and buffer C (0.01M Tris, pH 7.4 and 1% Nonidet P-40). The final pellets were solubilized with Laemmli buffer containing 100 mM dithiothreitol, boiled for 5 min, centrifuged in a microcentrifuge, and the supernatant was subjected to SDS-PAGE and Western blot analysis with anti-pTyr antibody.

[0082] **3. Western blot analysis**

[0083] Equal amount of hippocampal proteins were applied to SDS polyacrylamide gel. Electrotransfer of proteins from the gels to nitrocellulose paper (Schleicher & Schuell) was carried out for 1 hr at 100 V (constant) as described by Towbin et al. [Towbin H., Staehelin, J., Gordon, J. Electric transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. Proc. Natl. Acad. Sci. USA 76, 4350-4354, 1979]. The filter papers were preincubated for 1 hr at 23 C with PBS containing 0.1% Tween 20 and 3% bovine serum albumin and washed with PBS containing 0.1% Tween 20 three times for 10 min each. The blots were probed with pTyr antibodies for 1 hr at 23 C. The blots were then incubated with HRP-conjugated anti-rabbit IgG for 30 min and washed with PBS containing Tween 20 five times for 10 min each. The detection of immobilized specific antigens was carried out by ECL (NEN).

[0084] **3) Experimental results**

[0085] Since it has been suggested that tyrosine phosphorylation of the insulin receptors in hippocampus play an important role in spatial memory, we have tested whether AR extracts have any effect

on the tyrosine phosphorylation of hippocampal proteins. Tyrosine phosphorylation of a number of proteins with molecular sizes of ~180 kDa, 130 kDa, 95 kDa, 55 kDa and 42 kDa (Fig. 12). We further tested whether insulin receptor was tyrosine phosphorylated by the AR extracts. Insulin receptor phosphorylation was not detected under the basal condition whereas it was significantly stimulated by fraction 1 and fraction 2: The effect of fraction 1 was higher than that of fraction 2 (Fig. 13). In addition, the fraction 1, 2 and 4 of AR extracts has significantly stimulated ERK1 (44 kDa) and ERK2 (42 kDa) (Fig. 14).

[0086] Experiment 5. Ex vivo Cholinesterase assay

[0087] 1) Experimental methods

[0088] Male SD rats were dosed p.o. with vehicle or fractions of AR extract. The rats were decapitated after 90 min, brains rapidly removed, hippocampus and corpora striata dissected free, weighed and homogenized as described above. Cholinesterase activity was measured as described by Ellman et al [Ellman, G. L., Courtney, K. D., Andres, V., Featherstone, R. M. A new and rapid colorimetric determination of acetylcholinesterase activity. Biochem. Pharmacol. 7, 88-95.1961]. Briefly, 3 ml of buffer I (100 mM phosphate, pH 8.0), 0.2 ml of 75 mM acetylthiocholine iodide and 0.1 ml of buffered Ellman's reagent (DTNB 10 mM, NaHCO₃ 15 mM) were mixed and allowed to incubate for 10 min at 25°C. Then, 20 µl of enzyme sample was added and absorbance was measured at 30 sec intervals. The percent inhibition was calculated by comparison with the enzyme activity of the vehicle control group.

[0089] 2) Experimental results

[0090] Considering that cholinesterase inhibition has been shown to improve cognitive functions, we tested whether AR extracts could have some inhibitory effects on hippocampal cholinesterase activity. Administration of fraction 1, 2 or 4 caused an inhibition of hippocampal cholinesterase activity by 39%, 24% or 12% as compared to control, respectively (Fig. 15).

[0091] **Experiment 6. Oral Toxicity Test**

[0092] 1) Experimental methods

[0093] ICR mouse (20g) were housed for 1 week in a room with 50% relative humidity, 150-300 Lux, and controlled temperature (23 C) under 12 h light / dark cycle with free access to standard certified rodent diet and tap water. 25 mice were divided into 5 groups.

[0094] The Asiasari Radix extract (fraction 1) was dissolved in 0.1% Tween 80 and administered orally to 5 groups of the mice at the dose of 100 mg/Kg, 1,000 mg/Kg, 3,000 mg/Kg or 10,000mg/Kg, respectively. For the next 7 days, changes of general symptoms and death were observed. At day 7, all mice were killed and internal organs were examined.

[0095] 2) Experimental results

[0096] Administration of the Asiasari radix extract did not cause any changes in general symptoms and the appearance of internal organs. LD50 of the fraction 1 of the Asiasari Radix appeared as 3,400 mg/Kg.

[0097] **Preparation Example 1. Tablets**

[0098] Tablets are formulated by conventional manufacturing methods according to the following compositions.

[0099] 1-1. Tablets composition

Methanol extract of Asiasari Radix	500.0 mg
Lactose	500.0 mg
Talc	5.0 mg
Magnesium stearate	1.0 mg

[0100] 1-2. Tablets composition

Chloroform fraction of methanol extract of Asiasari Radix	50.0 mg
Lactose	50.0 mg
Talc	0.5 mg
Magnesium stearate	0.1 mg

[0101] 1-3. Tablets composition

Methanol fraction of methanol extract of Asiasari Radix	50.0 mg
Lactose	50.0 mg
Talc	0.5 mg
Magnesium stearate	0.1 mg

[0102] **Preparation Example 2. Capsules**

[0103] Capsules were prepared by the following methods according to the composition described below. Asiasari Radix extracts were sifted out and mixed with excipients and filled up in gelatin

capsules.

[0104] 2-1. Capsules composition

Methanol extract of Asiasari Radix	500.0 mg
Starch 1500	10.0 mg
Stearic acid magnesium BP	100 mg

[0105] 2-2. Capsules composition

Chloroform fraction of methanol extract of Asiasari Radix	50.0 mg
Starch 1500	1.0 mg
Stearic acid magnesium BP	10.0 mg

2-3. Capsules composition

Methanol fraction of methanol extract of Asiasari Radix	50.0 mg
Starch 1500	1.0 mg
Stearic acid magnesium BP	10.0 mg

[0106] Preparation Example 3. Syrups

Syrups were formulated by the following methods according to the composition described below. First of all, sugars were dissolved in purified water and paraoxybenzoate, paraoxypropylbenzoate and Asiasari Radix extracts were added. The resulting mixtures were dissolved at 60C and allowed to cool down. Finally, purified water was added to make final volume of 150 ml.

[0107] 3-1. Capsules composition

Methanol extract of Asiasari Radix	5.0 g
Sugar	95.1 g
Paraoxybenzoate	80.0 mg
Paraoxypropylbenzoate	16.0 mg
Purified water	to 150 ml

[0108] 3-2. Capsules composition

Chloroform fraction of methanol extract of Asiasari Radix	50.0 mg
Sugar	95.1 g
Paraoxybenzoate	80.0 mg
Paraoxypropylbenzoate	16.0 mg
Purified water	to 150 ml

[0109] 3-3. Capsules composition

Methanol fraction of methanol extract of Asiasari Radix	50.0 mg
Sugar	95.1 g
Paraoxybenzoate	80.0 mg
Paraoxypropylbenzoate	16.0 mg
Purified water	to 150 ml

[0110] Preparation Example 4. Solutions

[0111] The following compositions are formulated by conventional methods for solutions and the solutions are filled up in brown bottles.

[0112] 4-1. Solutions composition

Methanol extract of Asiasari Radix	500.0 mg
Isoglucose	20.0 g
Antioxidant	5.0 mg
Methylparabenzoic acid	2.0 mg
Distilled water	to 100.0 ml

[0113] 4-2. Solutions composition

Chloroform fraction of methanol

extract of Asiasari Radix	500.0 mg
Isoglucose	20.0 g
Antioxidant	5.0 mg
Methylparabenzoic acid	2.0 mg
Distilled water	to 100.0 ml

[0114] 4-3. Solutions composition

Methanol fraction of methanol	
extract of Asiasari Radix	500.0 mg
Isoglucose	20.0 g
Antioxidant	5.0 mg
Methylparabenzoic acid	2.0 mg
Distilled water	to 100.0 ml

[0115] Preparation Example 5. Powders

[0116] Powders were formulated as follows: the following components are mixed by conventional manufacturing methods for powders, filled into paper bags and completely sealed up.

[0117] 5-1. Powders composition

Methanol extract of Asiasari Radix	50.0 mg
Lactose	100.0 mg
Talc	5.0 mg

[0118] 5-2. Powders composition

Chloroform fraction of methanol extract of Asiasari Radix	50.0 mg
Lactose	100.0 mg
Talc	5.0 mg

[0119] 5-3. Powders composition

Methanol fraction of methanol extract of Asiasari Radix	50.0 mg
Lactose	100.0 mg
Talc	5.0 mg

[0120] Preparation Example 6. Injections

Injections were formulated by the following conventional manufacturing methods for injections according to the composition described below. The prepared injections are filled up in ampules of 2.0 ml volume and sterilized.

[0121] 6-1. Injections composition

Methanol extract of Asiasari Radix	50.0 mg
Antioxidant	1.0 mg

Tween 80	1.0 mg
Distilled water for injections	to 2.0 ml

[0122] 6-2. Injections composition

Chloroform fraction of methanol extract of Asiasari Radix	50.0 mg
Antioxidant	1.0 mg
Tween 80	1.0 mg
Distilled water for injections	to 2.0 ml

[0123] 6-3. Injections composition

Methanol fraction of methanol extract of Asiasari Radix	50.0 mg
Antioxidant	1.0 mg
Tween 80	1.0 mg
Distilled water for injections	to 2.0 ml

[0124] Effectiveness of the Invention

[0125] The composition containing Asiasari Radix extracts bring about not only prevention and treatment of neurodegenerative diseases caused by brain cell damages but also induction effect of memory improvement. It could be usefully utilized for the moderns who are exposed to dangerous brain damages due to various environmental stresses as well as for those who have decreased memory activity including dementia.

[0126] Additional modifications and improvements of the present

invention may also be apparent to those of ordinary skill in the art. Thus, the particular combination of parts described and illustrated herein is intended to represent only certain embodiments of the present invention, and is not intended to serve as limitations of alternative devices within the spirit and scope of the invention.

100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720 2721 2722 2723 2724 2725 2726 2727 2728 2729 2730 2731 2732 2733